

Idempotent biquadratics, Yang-Baxter maps and birational representations of Coxeter groups

James Atkinson

School of Mathematics and Statistics, the University of Sydney, NSW 2006, Australia

January 2013

Abstract: A transformation is obtained which completes the unification of quadrirational Yang-Baxter maps and known integrable multi-quadratic quad equations. A generalised discrete dynamics emerges by combining theory from these two classes of quad-graph models, it is encoded in a birational representation of a particular sequence of Coxeter groups. In this setting the usual quad-graph is associated with a subgroup of type BC_n , and is part of a larger and more symmetric ambient space. The model also defines, for instance, dynamics on a triangle-graph associated with a subgroup of type A_n , as well as finite degree-of-freedom dynamics, in the simplest cases associated with $\tilde{E}_6$ and $\tilde{E}_8$ affine subgroups. Underlying this structure is a class of biquadratic polynomials, that we call idempotent, which express the trisection of elliptic function periods algebraically via the addition law.

1. Introduction

The point of departure in this paper is the investigation of integrable quad equations. It will be demonstrated that the primary model in the multi-quadratic class that was discovered recently by the author in collaboration with Nieszporski [1], is related in a natural way to the primary model of the quadrirational Yang-Baxter maps that have been introduced earlier by Adler, Bobenko and Suris [2]. The existence of this relationship clarifies the position of the new model with respect to known quad-graph models and transformation theory in [3–15]. The subsequent focus of this article is on a particular repercussion of this connection, namely a new interpretation in terms of the *idempotent* class of biquadratic polynomials.

The idempotent biquadratic has its roots in the theory of elliptic functions; it is the formula for trisection of the periods expressed algebraically via the addition law. Interesting properties of this formula in part correspond to integrability of

the aforementioned quad-graph models, but go also beyond that, in particular leading to a natural generalisation of the quad-graph dynamics.

A practical way to understand the additional structure is in terms of the braid group representation emerging from the Yang-Baxter maps [16]. In the quadrirational case the braid-type generators are self-inverse, and the representation is only of the trivial part of the pure braid group, i.e., the symmetric group. But here we identify this as a subgroup of a larger ambient group. The precise characterisation of this larger group is one of the main technical achievements of this paper, it is a Coxeter group with unexpectedly rich structure (cf. Figure 10). For instance, in the sequence of groups formed by an increasing number of generators, the last finite case is the exceptional finite reflection group E_6 , and the associated lattice geometry coincides with the configuration of the 27 lines on a cubic surface. The general group structure bears closest resemblance to groups whose birational representation forms part of the basic framework in which the Painlevé equations and their generalisations may be understood [17–22].

We proceed as follows. Section 2 introduces the aforementioned quad-graph models and establishes the new relationship between them. The idempotent class of biquadratics is introduced in Section 3 with additional material given in Appendix A. Section 4 connects the quad-graph models with the idempotents via the notion of quadrirationality. A natural closure property of the idempotents, and an associated integrable triangle-graph dynamics, is established in Section 5. The nature of the relationship between the quad-graph and triangle-graph dynamics is a quite delicate matter, motivation behind a method to unify them is given by a local consideration of the problem in Section 6, and a constructive approach which completes the solution globally, based on vertex maps, is given in Section 7. The solution emerges in the form of the aforementioned birational group structure, which is interpreted from the point of view of lattice-geometry and dynamics in Section 8. Some consequences of the results here, and natural questions which emerge, are discussed in Section 9.

2. Preliminaries, F_I and $Q4^*$

The two relevant classes of models are the quadrirational Yang-Baxter maps and integrable multi-quadratic quad-equations. We recall the primary model from each class and connect them by a new Miura-type transformation.

Rational mappings with the Yang-Baxter property were constructed by Adler, Bobenko and Suris [2] on the basis of a simpler property termed quadrirationality. The primary model obtained was denoted F_I :

$$\begin{aligned}\hat{u} &= \alpha v \frac{(1-\beta)u + \beta - \alpha - (1-\alpha)v}{(1-\alpha)\beta u + (\alpha-\beta)uv - (1-\beta)\alpha v}, \\ \tilde{v} &= \beta u \frac{(1-\alpha)v + \alpha - \beta - (1-\beta)u}{(1-\beta)\alpha v + (\beta-\alpha)vu - (1-\alpha)\beta u}.\end{aligned}\tag{2.1}$$

It is natural to view such models as dynamical systems on a quad-graph [23–25], see Figure 1(a); variables assigned to edges are governed by the system (2.1) on each quad, whilst the essential parameters of the model, α and β , are associated with characteristics. This is unambiguous due to invariance of (2.1) under permutations $\hat{u} \leftrightarrow u$, $\tilde{v} \leftrightarrow v$ and $(u, \hat{u}, \beta) \leftrightarrow (v, \tilde{v}, \alpha)$, or in other words




Fig. 1. (a) Planar quad-graph, (b) quad with variables on edges, (c) quad with variables on vertices. Parameters are associated with the dashed lines, which represent characteristics.

the system respects the symmetry of the quad, see Figure 1(b). The rationality of the mapping plus the first two of these permutation symmetries is the essence of the quadrirationality.

A class of transformations connecting models with variables on edges, such as (2.1), with polynomial quad equations in which variables are on vertices, were introduced by Papageorgiou, Tongas and Veselov [10] and developed by Kassotakis and Nieszporski [12]. It is a new transformation in this class which is the point of departure in this article, it involves equations on edges of the graph which around a single quad take the form

$$\begin{aligned} B(u, x, \tilde{x}, \alpha) &= 0, & B(v, x, \hat{x}, \beta) &= 0, \\ B(\hat{u}, \hat{x}, \hat{\hat{x}}, \alpha) &= 0, & B(\tilde{v}, \tilde{x}, \hat{\tilde{x}}, \beta) &= 0, \end{aligned} \quad (2.2)$$

where B is the following polynomial

$$\begin{aligned} B(u, x, \tilde{x}, \alpha) &:= \\ (1 + cx)(1 + c\tilde{x})u^2 - [(1 + c^2)(1 + x\tilde{x})\alpha + 2c(x + \tilde{x})]u + (c + x)(c + \tilde{x})\alpha, \end{aligned} \quad (2.3)$$

and $c \in \mathbb{C} \setminus \{0, 1, -1, i, -i\}$ is a constant parameter. This system connects model (2.1) with the quad equation

$$\begin{aligned} (p - q)[(c^{-2}p - c^2q)(x\tilde{x} - \hat{x}\hat{\hat{x}})^2 - (c^{-2}q - c^2p)(x\hat{x} - \tilde{x}\hat{\tilde{x}})^2] \\ - (p - q)^2[(x + \hat{x})^2(1 + \tilde{x}^2\hat{x}^2) + (\tilde{x} + \hat{x})^2(1 + x^2\hat{\tilde{x}}^2)] \\ + [(x - \hat{\hat{x}})(\tilde{x} - \hat{x})(c^{-1} - cpq) - 2(p - q)(1 + x\tilde{x}\hat{x}\hat{\hat{x}})] \\ \times [(x - \hat{\hat{x}})(\tilde{x} - \hat{x})(c^{-1}pq - c) - 2(p - q)(x\hat{x} + \tilde{x}\hat{\tilde{x}})] = 0, \end{aligned} \quad (2.4)$$

where

$$p = m(\alpha), \quad q = m(\beta), \quad m := \alpha \mapsto \frac{2c^2 - (1 + c^2)\alpha}{c(1 + c^2)\alpha - 2c}, \quad (2.5)$$




Fig. 2. The Yang-Baxter property, global transfer of data along characteristics is independent of the order in which they cross. This is equivalent to the local condition that equations for variables on white nodes in terms of variables on black ones are the same for both of these quad graphs.

see Figure 1(c). The quad equation (2.4), denoted $Q4^*$, is quadratic in the dependent variable, but the basis of its original construction [1] guarantees existence of a rational reformulation. In fact the system on edges (2.2) is a way to introduce auxiliary edge variables, the (autonomous) rational reformulation in terms of these variables being exactly model (2.1). Typically such reformulation would involve both edge and vertex variables, so relations (2.2) are especially natural. It means system (2.2) is also a Miura-type transformation obtaining solutions of (2.4) from solutions of (2.1).

In fact these features can be explained in terms of the integrability of (2.1), namely the Yang-Baxter property, or cubic consistency, see Figure 2. The main observation is that

$$B(u, m(w), m(\tilde{w}), \alpha) = 0 \quad \Leftrightarrow \quad \tilde{w} = \gamma u \frac{(1 - \alpha)w + \alpha - \gamma - (1 - \gamma)u}{(1 - \gamma)\alpha u + (\gamma - \alpha)wu - (1 - \alpha)\gamma w}, \quad \gamma = \left(\frac{2c}{1 + c^2} \right)^2, \quad (2.6)$$

thus (2.2) is really nothing but the same underlying model (2.1). The system of all equations (2.1), (2.2), (2.4) is associated with the non-planar cube that, due to the consistency property, may be formed as the union of the two graphs in Figure 2 without breaking the well-posedness of the indicated initial value problem. This cube is shown in projected form in Figure 3 with variables and parameters added explicitly. It can be verified by calculation that as a consequence of imposing (2.1) on each face, the variables on the four edges through which the closed γ characteristic passes are related by (2.4) via the substitution

$$\begin{aligned} x &= m(w), \quad \tilde{x} = m(\tilde{w}), \quad \hat{x} = m(\hat{w}), \quad \hat{\hat{x}} = m(\hat{\hat{w}}), \\ p &= m(\alpha), \quad q = m(\beta), \quad c = m(\gamma). \end{aligned} \quad (2.7)$$

(Note that the generic form of (2.4) will be given later in this article, Möbius transformation m here and in (2.5) is characterised by its action $m : (0, 1, \infty) \mapsto (-c, 1/c, -1/c)$). Up to the same change of variables, the equations (2.2) are associated with the four quads through which the γ characteristic passes. Upon this observation the statement that (2.2) constitutes a Miura-type transformation between (2.1) and (2.4) can be identified with the statement of the Yang-Baxter property of (2.1) itself.

The theory of the simpler rational model (2.1) therefore encompasses the transformation (2.2) and the multi-quadratic model (2.4). In hindsight one can




Fig. 3. A quad-graph domain for (2.1). The four variables along the closed characteristic γ are related by the multi-quadratic quad equation (2.4) modulo the change of variables (2.7).

notice that the basis of construction of (2.1) and (2.4) are in fact similar. In the first instance it is the preservation of rationality under permutation of the variables of the model, and in the second it is existence of a rational reformulation.

However, there is one particular element from the theory of (2.4) which turns out to give new insight into (2.1). It is known [1] that the interpretation of B as Bäcklund transformation for (2.4) implies the following discriminant property,

$$\Delta[B(u, \mathbf{m}(w), \mathbf{m}(\tilde{w}), \alpha), u] = 0 \quad \Leftrightarrow$$

$$[(1 - \gamma)^{-1}(w - \gamma)(\tilde{w} - \gamma)(\alpha - \gamma) + \gamma(w + \tilde{w} + \alpha - \gamma)]^2 - 4\gamma w \tilde{w} \alpha = 0, \quad (2.8)$$

where the polynomial appearing in (2.8) is the Weierstrass-type symmetric tri-quadratic with discriminant polynomial $r(w) = w(w - 1)(w - \gamma)$. This characterises B as being amongst a particular class of polynomials that we call idempotent, as shown in Appendix A. Due to (2.6), the model (2.1) is also connected with this class of polynomials, and it is this connection which is the key observation we develop in this paper. The idempotent class has some remarkable features of independent interest, and the connection with (2.1) can be established with greater directness after some of those features are understood.

3. Idempotent biquadratic correspondences

We define and give basic properties of the relevant class of algebraic correspondences.

Symmetric biquadratic polynomials,

$$h(x, y) = c_0 + c_1(x + y) + c_2xy + c_3(x^2 + y^2) + c_4xy(x + y) + c_5x^2y^2, \quad (3.1)$$

$c_0, \dots, c_5 \in \mathbb{C}$, provided they don't decompose into a product involving one-variable polynomial factors, define dynamics on *orbits*, that is sequences of values

$$\dots, x_{-2}, x_{-1}, x_0, x_1, x_2, \dots \in \mathbb{C} \cup \{\infty\} \quad (3.2)$$

such that

$$h(x_n, y) = 0 \Leftrightarrow y \in \{x_{n-1}, x_{n+1}\}, \quad n \in \mathbb{Z}. \quad (3.3)$$

The two-valued dynamics defined by the biquadratic manifests only in the choice of direction along the orbit. The significance of the orbits, which are sequences defined uniquely up to orientation, is that they may be taken as a definition of elliptic functions in the discrete setting.

The self-composition of the correspondence defined by h yields another symmetric biquadratic polynomial, g , via the formula

$$\text{res}_2[h(x, y), h(y, z), y] = (x - z)^2 g(x, z), \quad (3.4)$$

where res_2 denotes the resultant of quadratic polynomials¹. The other factor appearing in (3.4), $(x - z)^2$, is due to the fact that h is symmetric. This notion of self-composition allows to define the idempotent class.

Definition 1. We refer to h (3.1) as idempotent if its self-composition g , defined by (3.4), is a non-zero scalar multiple of h .

Note that if h has a one-variable polynomial factor, or it drops in degree, then its self-composition vanishes identically. Such degenerate cases are excluded in Definition 1 by requiring the scalar multiple be non-zero. In particular, idempotent biquadratic polynomials have well-defined orbits.

The utility to be found in biquadratic polynomials from this class begins with the following list of complementary ways to characterise them.

Proposition 1. Assume the symmetric biquadratic polynomial h in (3.1) is not a scalar multiple of $(x - y)^2$, then the following are equivalent,

- (i) h is idempotent,
- (ii)* all orbits of h are three-periodic ($x_{n+3} = x_n$ in (3.2)),
- (iii)* there exists a non-constant three-periodic orbit of h ,
- (iv)* the coefficients of h satisfy the constraint $c_1 c_4 + c_3^2 = c_0 c_5 + c_2 c_3$,
- (v)* h satisfies the four-variable polynomial identity

$$(w - z)h(w, z)(x - y)h(x, y) + (w - y)h(w, y)(z - x)h(z, x) + (w - x)h(w, x)(y - z)h(y, z) = 0, \quad (3.5)$$

- (vi) h can be written in the form

$$h(x, y) = \frac{r_1(x)r_2(y) - r_2(x)r_1(y)}{x - y}, \quad (3.6)$$

where r_1, r_2 are cubic polynomials with no common roots.

Here the * indicates addition of the non-degeneracy condition on h that its self-composition, g in (3.4), does not vanish identically.

Proof. Deleting every other point from an orbit of h yields an orbit of its self-composition g defined by (3.4), considering this and the fact that orbits of g and h coincide due to (i), leads to (ii)*.

Existence of a non-constant orbit is a consequence of the assumption that h is not a scalar multiple of $(x - y)^2$, therefore (ii)* implies (iii)*.

¹ More precisely $\text{res}_2[p(x), q(x), x]$ is the determinant of the Sylvester matrix of p and q with the explicit assumption that both p and q are degree two polynomials, allowing a possibly zero coefficient of the second-degree monomial term.




Fig. 4. An orbit of an idempotent biquadratic: $h(x_i, y) = 0 \Leftrightarrow y \in \{x_j, x_k\}$, $\{i, j, k\} = \{0, 1, 2\}$.

To see that (iii)* implies (iv)* denote the elements of the three-periodic orbit mentioned in (iii)* by x_0 , x_1 and x_2 , at least two of which are distinct, and consider the resulting three equalities between polynomials in y :

$$\begin{aligned} h(x_0, y) &= (c_3 + c_4x_0 + c_5x_0^2)(y - x_1)(y - x_2), \\ h(x_1, y) &= (c_3 + c_4x_1 + c_5x_1^2)(y - x_2)(y - x_0), \\ h(x_2, y) &= (c_3 + c_4x_2 + c_5x_2^2)(y - x_0)(y - x_1). \end{aligned} \quad (3.7)$$

Substituting the explicit form of h (3.1) followed by elimination of x_0 , x_1 , x_2 yields the condition on the coefficients which appears in (iv)*.

Substitution of (3.1) directly shows that polynomial identity (3.5) is satisfied as a consequence of the condition on the coefficients appearing in (iv)*, so (iv)* implies (v)*.

Choosing w and z to be fixed constants such that $\eta^2 := (w - z)h(w, z) \neq 0$ it is clear that

$$\frac{1}{\eta}(w - x)h(w, x) = r_1(x), \quad \frac{1}{\eta}(z - x)h(z, x) = r_2(x), \quad (3.8)$$

are cubic polynomials in x , and writing (3.5) in terms of these one obtains (3.6). Furthermore r_1, r_2 have no common roots due to the non-degeneracy assumption on h , therefore (v)* implies (vi).

Assuming h is of the form (3.6) specified in (vi), a direct calculation shows that

$$\text{res}_2[h(x, y), h(y, z), y] = (x - z)^2 \mu h(x, z), \quad \mu := \text{res}_3[r_1(y), r_2(y), y]. \quad (3.9)$$

The scalar multiple μ appearing in (3.9) is non-zero if and only if r_1, r_2 in (3.6) have no common roots (res_3 excluding also the possibility of ∞ being a common root, which corresponds to a drop in degree of both polynomials), therefore (vi) implies (i).

Equivalences (ii)* and (iii)* connect the idempotent class of biquadratics with the trisection of the elliptic function periods. The three-periodic orbits will play the central role in what follows. They are specified completely by the un-ordered triplet of participating points, orbits being determined only up to orientation. Therefore within this class of biquadratics the un-ordered triplet itself may be referred to as the orbit, and, to reflect the symmetry, it is natural to think of points in an orbit as being arranged in a triangle as in Figure 4.

We remark that the idempotents occupy a similar special position in relation to the biquadratic correspondences as do Möbius involutions to the group of

Möbius transformations. This was observed in [26] based on the similarity of characteristic features of Proposition 1 to characteristics of the Möbius involutions, but can also be seen directly by considering formula (3.6) in the degenerate case when r_1 and r_2 are degree two polynomials.

Expression (3.6) makes connection with Bezout's formula for the resultant of polynomials r_1 and r_2 (see [27]), specifically the resultant is the determinant of the coefficient matrix of (3.6). Its appearance here is interesting, but perhaps not completely surprising given the role played by the resultant in Definition 1.

As a final remark, we note that the pattern of terms and variables in (3.5) is very similar to that of the three-term sigma-function identity (see p. 390 of [28]), or Riemann relation, though an explanation for this is not so clear.

The characterising polynomial identity (3.5) leads directly to the following constructive existence-and-uniqueness result for the idempotent biquadratics.

Lemma 1. *Given a pair of disjoint triplets taken from $\mathbb{C} \cup \{\infty\}$*

$$x_0, x_1, x_2, \quad y_0, y_1, y_2, \quad (3.10)$$

there exists an idempotent biquadratic, which is unique up to a scalar multiple, for which these are orbits. It may be expressed via the formula (3.6) by choosing

$$r_1(x) = (x - x_0)(x - x_1)(x - x_2), \quad r_2(x) = (x - y_0)(x - y_1)(x - y_2). \quad (3.11)$$

Proof. For existence it is sufficient to check that the roots of the polynomials r_1 and r_2 give two particular orbits of h in (3.6). Conversely, supposing (3.10) are the orbits of a given idempotent biquadratic polynomial h , choose $w = x_0$ and $z = y_0$ in the identity characterising such polynomials (3.5). By inspection it can be seen that the polynomials r_1, r_2 appearing in (3.6), (3.8), which is just a re-arrangement of (3.5), vanish on the orbits (3.10) due to the choice of w and z . In other words, any such biquadratic h coincides with (3.6), (3.11) up to a scalar multiple.

Thus two orbits can be chosen freely, which then determine an idempotent biquadratic polynomial uniquely up to a scalar multiple. Clearly three orbits cannot be chosen freely.

4. Rationality of the three-orbit constraint

We connect the previously defined class of algebraic correspondences with the scalar quadrirational dynamical systems, this emerges directly by looking more closely at the three-orbit constraint.

The following form of this constraint is a straightforward consequence of Lemma 1.

Proposition 2. *Denote three elements in the vector space of cubic polynomials by r_1, r_2 and r_3 . The roots of these three polynomials are three distinct orbits of an idempotent biquadratic correspondence if and only if there are no pair-wise common roots and they are linearly dependent:*

$$\exists \gamma_1, \gamma_2, \gamma_3 \in \mathbb{C} \setminus \{0\} \text{ such that } \gamma_1 r_1(x) + \gamma_2 r_2(x) + \gamma_3 r_3(x) = 0 \quad \forall x. \quad (4.1)$$




Fig. 5. Quadrirationality: variables on any pair of adjacent edges are determined rationally from the variables on the opposing edges.

In terms of the roots of r_1, r_2, r_3 , i.e., directly in terms of the points of the three orbits, say

$$x_0, x_1, x_2, \quad y_0, y_1, y_2, \quad z_0, z_1, z_2, \quad (4.2)$$

the constraint leaves seven degrees of freedom amongst the nine variables. An important feature is its symmetry, by which we mean invariance under permutation of the three orbits, as well as permutation of the three points of any one of those orbits. The main reason for drawing attention to this symmetry is that, although transparent on the level of Proposition 2, it is obscured when writing the constraint in a more explicit form.

One possible explicit form appears by specifying all variables in (4.2) except two taken from the same orbit. For instance, z_1 and z_2 are determined in terms of the remaining variables by a quadratic equation, $h(z_0, z) = 0 \Leftrightarrow z \in \{z_1, z_2\}$, with h given by (3.6), (3.11). However, if two variables are chosen from distinct orbits, then they are determined rationally. To see this consider how the biquadratic (3.6), (3.11) depends on all data

$$h(x, y; x_0, x_1, x_2; y_0, y_1, y_2) = (x - y)^{-1} [(x - x_0)(x - x_1)(x - x_2)(y - y_0)(y - y_1)(y - y_2) - (y - x_0)(y - x_1)(y - x_2)(x - y_0)(x - y_1)(x - y_2)]. \quad (4.3)$$

The dependence is quadratic in x and y , but in the remaining variables h has degree one. Writing the constraint on the three orbits (4.2) in the form

$$h(x_0, x_1; z_0, z_1, z_2; y_0, y_1, y_2) = 0, \quad h(y_0, y_1; z_0, z_1, z_2; x_0, x_1, x_2) = 0, \quad (4.4)$$

it is clear from the degree of the polynomials that y_2 and x_2 are determined rationally from the remaining variables.

Consider now combining the rationality of the constraint with its permutation symmetries $x_1 \leftrightarrow x_2$ and $y_1 \leftrightarrow y_2$. It is clear that with x_0, y_0, z_0, z_1 and z_2 fixed, the constraint on the remaining four variables x_1, x_2, y_1 and y_2 , visualised by assigning variables to edges of a quad as in Figure 5, determines any pair of adjacent variables rationally from those on the opposing edges. This system therefore falls within the *quadrirational* class introduced in [2]. An exhaustive list of five scalar quadrirational models was found, and denoted $F_I, \dots, F_V$. The primary model, F_I , was written in the introduction (2.1), it is exactly the condition required for existence of an idempotent biquadratic polynomial with the three particular orbits

$$u, \hat{u}, \beta, \quad v, \tilde{v}, \alpha, \quad 0, 1, \infty. \quad (4.5)$$

Notice that the third orbit is constant. When two or all three points of the constant orbit coincide, the associated quadrirational model is equivalent to F_{II} or F_{IV} respectively. The remaining models, F_{III} and F_V , are obtained by limiting procedures that erode the connection to the idempotent biquadratics and are not considered here.

The dynamics defined by the quadrirational models F_I , F_{II} and F_{IV} are therefore associated with a family of idempotent correspondences that have a common fixed orbit, the distinction between the three models being the number of points in that orbit. This distinction is not important here, where possible we consider the generic form of these models (4.4) in which variables z_0 , z_1 and z_2 of the third orbit are assumed fixed, but are left unspecified.

The next step we take in this paper is the introduction of a second class of integrable dynamics, distinct from the quad-graph, which emerges from elementary considerations of this family of algebraic correspondences.

5. Oriented tetrahedral consistency

We establish a closure/associativity property in the family of idempotent biquadratic correspondences that share a common orbit. This can be seen as a consistency feature of the rational three-orbit constraint (4.4). It allows to interpret it as a discrete integrable dynamical system whose domain is the (edge-) oriented n -simplex.

The relevant closure property may be stated as follows.

Proposition 3. *Denote twelve distinct elements taken from $\mathbb{C} \cup \{\infty\}$ by*

$$x_{ij}, \quad i, j \in \{1, 2, 3, 4\}, \quad |\{i, j\}| = 2, \quad (5.1)$$

and take

$$x_{ij}, x_{jk}, x_{ki}, \quad x_{ji}, x_{kj}, x_{ik}, \quad i, j, k \in \{1, 2, 3, 4\}, \quad |\{i, j, k\}| = 3, \quad (5.2)$$

to be pairs of orbits that define (cf. Lemma 1) four idempotent biquadratic polynomials. If there exists an orbit common amongst any three of these, then it is common amongst all four.

The geometric object we associate with this arrangement of variables is the tetrahedron with oriented edges. Indices are associated with vertices, variables with edges, and orbits with faces, see Figures 6(a) and 6(b).

More algebraic intuition is offered by interpreting this property as associativity. Consider the binary operation

$$(x_0, y_0) \cdot (x_1, y_1) := (y_2, x_2) \quad (5.3)$$

defined by system (4.4) when z_0 , z_1 and z_2 are considered fixed. The commutativity of this operation is clear from the permutation symmetry of the defining system, the associativity can be verified directly by calculation. This is equivalent to verifying Proposition 3, which can be seen as follows.

The proposition states that if the three-orbit constraint (4.4) is satisfied for three of the idempotents, then it is satisfied for the fourth. Taking the common




Fig. 6. (a) Tetrahedron, (b) one edge and two variables, one variable is assigned to each orientation of the edge, (c) a planar triangle tessellation.

third orbit to be z_0, z_1, z_2 , three of the constraints written in terms of the above binary operation are as follows

$$\begin{aligned} (x_{14}, x_{41}) &= (x_{13}, x_{31}) \cdot (x_{34}, x_{43}), \\ (x_{24}, x_{42}) &= (x_{21}, x_{12}) \cdot (x_{14}, x_{41}), \\ (x_{23}, x_{32}) &= (x_{21}, x_{12}) \cdot (x_{13}, x_{31}). \end{aligned} \quad (5.4)$$

Assuming these hold one sees by substitution that the fourth constraint,

$$(x_{24}, x_{42}) = (x_{23}, x_{32}) \cdot (x_{34}, x_{43}), \quad (5.5)$$

is equivalent to the associativity condition

$$(x_{21}, x_{12}) \cdot [(x_{13}, x_{31}) \cdot (x_{34}, x_{43})] = [(x_{21}, x_{12}) \cdot (x_{13}, x_{31})] \cdot (x_{34}, x_{43}). \quad (5.6)$$

That the equations represented by (5.5) are a consequence of those represented by (5.4) means this is a consistency property similar in spirit to the known integrability feature of the quadrirational models, namely the consistency on a cube, or Yang-Baxter property. Here equations on three faces of the tetrahedron (5.4) imply the equation on the fourth face (5.5).

The natural extension of this domain is the n -simplex. Coordinatization is immediate by replacing the range of indices $\{1, 2, 3, 4\}$, appearing the definition of variables (5.1) and orbits (5.2) of the 3-simplex, with the set $\{1, \dots, n+1\}$. Tessellation by triangles of any two-dimensional surface (see Figure 6(c)) can be considered as a sub-case of this domain, which is clear because the n -simplex can be recovered by adding edges until all pairs of vertices are joined.

Models with such tetrahedral consistency can be directly integrated. Specifically the general solution of a given initial value problem can be written down in terms of the binary operation. Natural initial data for the oriented n -simplex domain is the set of pairs of variables associated with all edges connected to some vertex, for instance the n pairs (x_{ij}, x_{ji}) , $j \in \{1, \dots, n+1\} \setminus \{i\}$ connected

z_0, z_1, z_2	$\varphi(a, b)$	$\varphi(a, b) \cdot \varphi(c, d)$
$0, 1, \infty$	$(a(1-b)/(a-b), (1-b)/(a-b))$	$\varphi(ac, bd)$
$0, \infty, \infty$	$(a/(b-1), ab/(b-1))$	$\varphi(a+c, bd)$
∞, ∞, ∞	$(a/b+b, a/b-b)$	$\varphi(a+c, b+d)$

Table 1. Decomposing the binary operation into addition and multiplication.

to vertex i . In this case the general solution in terms of the initial data is given by $(x_{jk}, x_{kj}) = (x_{ji}, x_{ij}) \cdot (x_{ik}, x_{ki})$ for any $j, k \in \{1, \dots, n+1\} \setminus \{i\}$.

The binary operation (5.3) defined by (4.4) is of itself very simple, it can be decomposed into addition and multiplication. Substitutions φ achieving this are listed in Table 1 for three canonical choices of the fixed third orbit.

It is natural to ask what is the relationship between the dynamics observed here and the usual quad-graph dynamics. Answering this question is in fact the main technical focus of the remainder of this paper. We begin in the following section with a local consideration of this problem, establishing a combinatorial relationship between the cubic consistency and edge-oriented tetrahedral consistency properties.

6. 5-simplex consistency

Through a symmetrization procedure we introduce a particular domain for model (4.4) which is combinatorially equivalent to the 5-simplex. On this domain the usual cubic consistency (established in [2] for all quadrirational models) and the oriented-tetrahedron consistency (introduced in Section 5) are exhibited simultaneously as sub-cases of a stronger consistency property.

6.1. Completeness based on symmetry. It is useful to broaden the notion of domain for (4.4) to a more combinatorial setting. Considered with the orbit z_0, z_1, z_2 fixed, the system (4.4) has the symmetry of a pair of disjoint triangles, therefore in general the domain can be a set of variables arranged into *triplet-pairs*, where:

Definition 2. A *triplet-pair* is a disjoint pair of sets, each set containing three variables.

To remove degeneracy we assume connectedness, i.e., that any pair of variables in the domain are connected through some sequence of triplet-pairs.

The domain appearing in Proposition 3 contains in total four triplet-pairs, whilst each variable participates in exactly two of them. A domain with this kind of regularity will be termed a *pair-configuration*.

Definition 3. By a *pair-configuration* we mean a set of variables arranged into triplet-pairs such that each variable participates in the same number of triplet-pairs.

It means that throughout the domain, like within the equation itself, all variables are on an equal footing. We remark that neglecting the pair associations between triplets recovers an abstract configuration in the usual sense. Also we




Fig. 7. The first diagram is the 5-simplex, fifteen variables are assigned to the fifteen edges, and ten triplet-pairs are assigned to the ten vertex-disjoint pairs of triangles, or *polar* triangles, a typical example of which is illustrated in the second diagram.

deal exclusively with triplets here, but the more general definition would involve n -tuples of variables.

By a symmetry of a pair-configuration we mean a permutation of the variables which also permutes the triplet-pairs. The group of symmetries of the simplest pair-configuration, a single triplet-pair (3.10), contains elements that permute each triplet, and also those that swap the two triplets. This group has order $3! \times 3! \times 2 = 72$. On the other hand, the oriented tetrahedral pair-configuration has just $2 \times 4! = 48$ symmetries, corresponding to the tetrahedral symmetry plus the overall reversal of orientation. By counting, it is therefore clear that not every symmetry of each participating triplet-pair extends to a symmetry of the full tetrahedral pair-configuration. For a pair-configuration which is a domain of our model we adopt the principle that this is a deficiency. To overcome it we seek to extend the configuration, such extension will be termed symmetry-complete.

Definition 4. A pair-configuration is symmetry-complete when every symmetry of every triplet-pair is the restriction of a symmetry of the whole configuration.

It is natural to require the extension be minimal, that is to ask for the *smallest* symmetry-complete pair-configuration that contains the original. The question of whether a finite completion of this kind exists is purely combinatorial. For the pair-configuration of Proposition 3 the answer is affirmative.

Proposition 4. The smallest symmetry-complete extension of the edge-oriented tetrahedral pair-configuration (cf. Proposition 3) consists of fifteen variables

$$x_{ij} = x_{ji}, \quad i, j \in \{1, 2, 3, 4, 5, 6\}, \quad |\{i, j\}| = 2, \quad (6.1)$$

arranged into the ten triplet-pairs

$$x_{ij}, x_{jk}, x_{ki}, \quad x_{lm}, x_{mn}, x_{nl}, \quad \{i, j, k, l, m, n\} = \{1, 2, 3, 4, 5, 6\}. \quad (6.2)$$

This extended configuration has the combinatorics of the 5-simplex, see Figure 7. The indices are assigned to vertices of the figure, the variables to edges and the triplet-pairs to pairs of vertex-disjoint (or polar) triangles.

Proof (of Proposition 4). That the new configuration is symmetry-complete in the sense of Definition 4 can be seen from the figure, the symmetry group of the 5-simplex is S_6 which acts naturally on its vertices. By inspection this group

acts transitively on the triangle-pairs, whilst the sub-group that stabilizes any particular triangle-pair is the complete symmetry group of that pair.

To see that this configuration is minimal consider the intersection between two distinct triplet-pairs. In the case of the tetrahedral configuration the intersection is of a particular nature, triplet-pairs intersect on two variables - one from each triplet. For a hypothetical symmetry-complete extension of this configuration it is necessary that for any participating triplet-pair and any two variables chosen one from each of its triplets, that there exist another triplet-pair intersecting the first on those variables. For the 5-simplex pair-configuration described, a triplet-pair intersecting in this way exists and is in fact *unique*, and furthermore *all* intersections are of this kind. The extended configuration therefore cannot be any smaller, for if it were it would omit some of the necessary intersections.

6.2. The stronger consistency property. The symmetrization procedure thus extends the tetrahedral domain to one which respects the symmetry of the model. It is a remarkable additional feature of (4.4) that the well-posedness of the initial value problem is also preserved in this extension. In order to describe this situation precisely the following notion is useful.

Definition 5. *A duad-syntheme of the 5-simplex is a triplet of vertex-disjoint edges.*

This terminology follows Sylvester [29], a more precise usage would be *edges associated with a duad-syntheme*. The *duads* here are un-ordered pairs from the set $\{1, \dots, 6\}$, and a duad-syntheme is a partition of this set into duads. In the same terminology a pair of vertex-disjoint triangles, to which we have previously assigned a triplet-pair, would be called a *triad-syntheme*. There are fifteen distinct duad-synthemes in total, the associated variables are given explicitly as follows

$$x_{ij}, \quad x_{kl}, \quad x_{mn}, \quad \{i, j, k, l, m, n\} = \{1, 2, 3, 4, 5, 6\}. \quad (6.3)$$

The notion of syntheme allows concise statement of the consistency feature.

Proposition 5. *Consider model (4.4) on the 5-simplex domain (i.e., the domain described in Proposition 4). Generic initial data on any pair of disjoint duad-synthemes determines a unique solution.*

This can be verified directly by calculation. Only a single calculation is necessary because any pair of disjoint duad-synthemes can be transformed to any other pair by the action of the 5-simplex symmetry group. Note that this does not exhaust the initial data configurations that determine a unique solution.

The consistency property may be understood more clearly through the following decomposition which again involves the duad-synthemes, but in a different manner. Distinguish any one of the duad-synthemes and partition the triplet-pairs of the 5-simplex pair-configuration into two sets according to whether or not they intersect it. There are four triplet-pairs disjoint from the distinguished syntheme, they lie in the oriented-tetrahedron pair-configuration of Proposition 3. The remaining six triplet-pairs are in natural correspondence with the six faces of a cube with characteristics. To illustrate the association we have labelled the edges of the 5-simplex in Figure 8 according to the notation of the cube in Figure 3. The parameters, α , β and γ lie on a duad-syntheme which




Fig. 8. The 5-simplex with variables assigned to edges.

induces the following partition of triplet-pairs:

$$\begin{array}{cc|cc}
 u, \bar{v}, \hat{w} & w, \tilde{v}, \bar{u} & \alpha, v, \tilde{v} & \beta, u, \hat{u} \\
 v, \tilde{w}, \bar{u} & u, \hat{w}, \tilde{v} & \beta, w, \hat{w} & \gamma, v, \bar{v} \\
 w, \hat{u}, \tilde{v} & v, \bar{u}, \hat{w} & \gamma, u, \bar{u} & \alpha, w, \tilde{w} \\
 \hat{u}, \bar{v}, \tilde{w} & \bar{u}, \tilde{v}, \hat{w} & \alpha, \bar{v}, \tilde{v} & \beta, \bar{u}, \hat{u} \\
 & & \beta, \tilde{w}, \hat{w} & \gamma, \tilde{v}, \tilde{v} \\
 & & \gamma, \hat{u}, \bar{u} & \alpha, \hat{w}, \hat{w}
 \end{array} \quad (6.4)$$

One can verify the oriented tetrahedron pair-configuration on the left, whilst the six triplet-pairs on the right can be identified with faces of the cube domain of Figure 3. Consistency on the cube is usually described in terms of initial data α, β, γ and u, v, w . This initial data allows to see equations on the left in (6.4) as a consequence of equations on the right. But there are fifteen ways to make such a partition, corresponding to the fifteen distinct duad-syntheses. There is therefore a harmony between the cube and tetrahedral consistencies, and it is natural to see the 5-simplex consistency as a stronger property of (4.4) which contains both as sub-cases.

6.3. The multi-quadratic model $Q4^*$. In this setting, which generalises the cube, the multi-quadratic model $Q4^*$ relates variables on the edges shown in Figure 9. This equation, returning to the notation of Proposition 4 (which reflects better the symmetry, corresponding to permutations of indices $1, \dots, 6$ of the variables x_{ij}), takes the following generic form

$$\begin{aligned}
 \psi(z_0, z_1, z_2) + \psi(z_1, z_2, z_0) + \psi(z_2, z_0, z_1) + \\
 \psi(z_2, z_1, z_0) + \psi(z_0, z_2, z_1) + \psi(z_1, z_0, z_2) = 0, \quad (6.5)
 \end{aligned}$$




Fig. 9. An edge of the 5-simplex and its polar tetrahedron.

where

$$\begin{aligned} \psi(z_0, z_1, z_2) &= (x_{12} - z_2)(z_0 - z_1) \operatorname{cr}(x_{34}, z_1, x_{56}, z_2) \operatorname{cr}(x_{34}, z_2, x_{56}, z_0) \\ &\quad \times (\operatorname{cr}(x_{45}, x_{12}, x_{36}, z_2) - \operatorname{cr}(x_{46}, z_1, x_{35}, z_0)) \\ &\quad \times (\operatorname{cr}(x_{45}, z_1, x_{36}, z_0) - \operatorname{cr}(x_{46}, x_{12}, x_{35}, z_2)), \end{aligned} \quad (6.6)$$

and cr denotes the cross-ratio

$$\operatorname{cr}(a, b, c, d) := \frac{(a - b)(c - d)}{(a - c)(b - d)}. \quad (6.7)$$

It involves the variable x_{12} , which is singled out, and variables on the edges of the tetrahedron formed by the remaining vertices 3, 4, 5, 6. The tetrahedral permutation symmetry of this model can be verified, though it is not obvious from (6.5), or from the expression (2.4), which corresponds to the particular case of (6.5) when $x_{12} = c$ and $\{z_0, z_1, z_2\} = \{-c, 1/c, -1/c\}$. The 5-simplex domain carries fifteen copies of this equation, corresponding to the fifteen edge-tetrahedron pairs, cf. Figure 9.

6.4. A point-line configuration equivalent to the 5-simplex pair-configuration. We show a natural way in which the 5-simplex pair-configuration can be viewed as a point-line configuration in a three-dimensional projective space.

Take the fifteen variables to be points and the set of all triplets from the triplet-pairs (6.2) and the duad-syntheses (6.3) to be lines (sets containing three points), so there are 35 lines in total. It can be verified that these are the 15 points and 35 lines of $PG(3, 2)$, i.e., the projective geometry in three dimensions defined over the field of characteristic two. The subset of 20 lines (6.2), whose union contains the entire set of 15 points, is the configuration of interest. Each point is on four lines and each line contains three points, it is a $(15_4 20_3)$ configuration. In $PG(3, 2)$ each line is contained in three planes, and within this particular configuration there is a unique other line, skew to the first, which passes through all three of those planes. This relationship is reciprocal, and it recovers the pair associations of the pair-configuration. Whereas the system (4.4) lies on these pairs of skew lines, the equation (6.5) lies on planes. The complement of this configuration in $PG(3, 2)$, i.e., the set of lines corresponding to the duad-syntheses (6.3), is known as the Cremona-Richmond configuration [30].

7. The vertex-map group

It is natural to ask what is the more general domain for models such as (4.4) which have the stronger 5-simplex consistency property. Formulation in terms of vertex maps allows group analysis to be applied to this question. To introduce the vertex maps we re-visit the oriented n -simplex and n -dimensional hypercube domains natural for models with the oriented-tetrahedral and cubic consistency properties respectively.

7.1. Vertex-maps of the oriented n -simplex. The natural initial-value problem for system (4.4) on the oriented n -simplex domain (cf. Section 5) suggests the following elementary construction.

Choose some vertex of the oriented n -simplex and denote the pairs of variables associated with its n connected edges by

$$\begin{bmatrix} x_1, y_1 \\ \vdots \\ x_n, y_n \end{bmatrix}, \quad (7.1)$$

the order in which edges are chosen is free, but we have chosen x_i to denote the departing variable and y_i to denote the arriving one (cf. Figure 6(b)). From such initial data the system (4.4) determines rationally all remaining edge variables of the n -simplex, and in particular the data similar to (7.1) associated with any other vertex. In particular, on the new vertex connected to the initial vertex by initial edge j , this data is obtained by a mapping that we denote τ_j ,

$$\tau_j := \begin{bmatrix} x_1, y_1 \\ \vdots \\ x_n, y_n \end{bmatrix} \mapsto \begin{bmatrix} \bar{x}_1, \bar{y}_1 \\ \vdots \\ \bar{x}_n, \bar{y}_n \end{bmatrix}, \quad (7.2)$$

where

$$\begin{aligned} (\bar{x}_i, \bar{y}_i) &= (y_j, x_j) \cdot (x_i, y_i), \quad i \in \{1, \dots, n\} \setminus \{j\}, \\ (\bar{x}_j, \bar{y}_j) &= (y_j, x_j). \end{aligned} \quad (7.3)$$

This mapping is expressed here in terms of the previously introduced binary operation (5.3) defined by (4.4).

The mappings $\tau_1, \dots, \tau_n$ generate a group which we call the n -simplex vertex-map group. It is isomorphic to the full symmetry group of the n -simplex, A_n . The generators satisfy the relations

$$\begin{aligned} \tau_i^2 &= \text{id}, \\ (\tau_i \tau_j)^3 &= \text{id}, \quad |\{i, j\}| = 2, \\ (\tau_i \tau_j \tau_k)^2 &= \text{id}, \quad |\{i, j, k\}| = 3, \end{aligned} \quad (7.4)$$

which, it can be verified, constitutes a (non-standard) presentation for this group. The order of the group is $(n+1)!$. If we consider the vertex data (7.1) only up to the ordering of connected edges ($n!$ permutations), then the natural group action corresponds to permutations of the $n+1$ vertices.

The relations (7.4) encode the oriented-triangle permutation symmetries and oriented-tetrahedral consistency property of (4.4).

7.2. Vertex-maps of the n -dimensional hypercube. The hypercube is the generic quad-graph domain for models with the cubic consistency property, containing for instance the planar quad-graph (with no self-intersecting characteristics) by a procedure of restriction. This domain also allows to define the vertex maps.

Consider the following regular initial-value-problem. Denote by $y_1, \dots, y_n$ the parameters associated with the n characteristics of the n -cube. Then select a vertex and denote by $x_1, \dots, x_n$ the variables on edges connected to that vertex, with ordering chosen so that the edge associated with variable x_i intersects the characteristic associated with parameter y_i . Arrange the n variable-parameter pairs in the array (7.1), the overall ordering of pairs may be chosen freely.

From such initial data the system (4.4) determines rationally all remaining edge variables, and in particular the similar data associated with any other vertex of the n -cube. On the vertex at the other end of initial edge j the similar data is obtained through a mapping which we denote by σ_j :

$$\sigma_j := \begin{bmatrix} x_1, y_1 \\ \vdots \\ x_n, y_n \end{bmatrix} \mapsto \begin{bmatrix} \bar{x}_1, y_1 \\ \vdots \\ \bar{x}_n, y_n \end{bmatrix}, \quad (7.5)$$

where $\bar{x}_1, \dots, \bar{x}_n$ are determined in terms of polynomial h (4.3) by equations

$$\begin{aligned} h(x_j, y_i; z_0, z_1, z_2; x_i, y_j, \bar{x}_i) &= 0, \quad i \in \{1, \dots, n\} \setminus \{j\}, \\ \bar{x}_j &= x_j. \end{aligned} \quad (7.6)$$

The mappings $\sigma_1, \dots, \sigma_n$ correspond to reflections normal to each axis of the n -cube, in particular they satisfy the relations

$$\begin{aligned} \sigma_i^2 &= \text{id}, \\ (\sigma_i \sigma_j)^2 &= \text{id}, \quad |\{i, j\}| = 2, \end{aligned} \quad (7.7)$$

which encode the quad-symmetry and cubic consistency (or Yang-Baxter property) of (4.4). These mappings generate a group of order 2^n and preserve the ordering of characteristics. Combined with pure permutation of the characteristics, i.e., the $n!$ maps that permute the rows of (7.1), they generate the full symmetry group of the n -dimensional hypercube, BC_n of order $2^n n!$. We call this the vertex-map group of the n -cube.

It is possible here to make contact with the well-known braid-group formulation of the Yang-Baxter property. It corresponds to a similar construction to the one above, but a path-type initial value problem for the n -cube. Consider again specifying the parameters on characteristics but now initial data on a sequence of n edges connecting two (hyper-) opposite vertices of the n -cube. We write this data as an array of n edge-variables each paired with the parameter associated with the characteristic that edge crosses, so the data takes the same form (7.1). The group of mappings that take one such initial data path to another are generated by those that take the simple form

$$s_j := \begin{bmatrix} x_1, y_1 \\ \vdots \\ x_n, y_n \end{bmatrix} \mapsto \begin{bmatrix} \bar{x}_1, \bar{y}_1 \\ \vdots \\ \bar{x}_n, \bar{y}_n \end{bmatrix}, \quad (7.8)$$

where $j \in \{1, \dots, n-1\}$ and

$$\begin{aligned} \bar{x}_i &= x_i, \quad \bar{y}_i = y_i, \quad i \in \{1, \dots, n\} \setminus \{j, j+1\}, \\ h(x_{j+1}, y_{j+1}; z_0, z_1, z_2; x_j, y_j, \bar{x}_j) &= 0, \quad \bar{y}_j = y_{j+1}, \\ h(x_j, y_j; z_0, z_1, z_2; x_{j+1}, y_{j+1}, \bar{x}_{j+1}) &= 0, \quad \bar{y}_{j+1} = y_j. \end{aligned} \quad (7.9)$$

They correspond to deforming the path by flipping edges across a single quad, in other words interchanging adjacent characteristics. The characteristics correspond to braids and the Yang-Baxter property is encoded in the first of the braid relations

$$\begin{aligned} s_i s_j s_i &= s_j s_i s_j, & |i - j| &= 1, \\ s_i s_j &= s_j s_i, & |i - j| &> 1. \end{aligned} \quad (7.10)$$

Here $s_i^2 = \text{id}$, so for these systems the braid relations lead only to the symmetric group. The intended extension beyond this group (which means extension beyond the hypercube) is initially less transparent in terms of the braid-type mappings. On the other hand, the hypercube geometry makes it straightforward to construct a transformation ρ that takes the vertex-type initial data to the path-type initial data. Denote by ϕ the column vector (7.1) and define ρ by the expression

$$\rho := \phi \mapsto \text{diagonal}([\phi, \sigma_1(\phi), [\sigma_1 \sigma_2](\phi), \dots, [\sigma_1 \cdots \sigma_{n-1}](\phi)]). \quad (7.11)$$

The inverse is obtained similarly,

$$\rho^{-1}(\phi) = \text{first-row}([\phi, s_1(\phi), [s_1 s_2](\phi), \dots, [s_1 \cdots s_{n-1}](\phi)]^T). \quad (7.12)$$

Note that arrays in (7.11) and (7.12) are to be understood as $n \times n$ square matrices whose entries are pairs of variables such as (x_1, y_1) etc. From this explicit form for ρ it is possible to verify, for instance, that the transformed braid-type maps,

$$\rho^{-1} s_1 \rho, \dots, \rho^{-1} s_{n-1} \rho, \quad (7.13)$$

are standard generators of the pure permutation part of the hypercube vertex-map group.

Note that from here it is possible to make contact also with the transfer dynamics of Veselovs monodromy maps [31]. In terms of the lattice geometry it corresponds to tessellation of $\mathbb{Z}^n$ by hypercubes, and the monodromy maps are identified with dynamics of the periodic, or finite-gap, reduction (details of this were established in the context of more general intertwining Yang-Baxter maps by Kouloukas and Papageorgiou [32]).

7.3. Characterisation of the complete vertex-map group. The previous two subsections establish transformation groups that act naturally on *vertex-type* initial data of the n -dimensional edge-oriented simplex and hypercube respectively. Combining both sets of generators in the case $n = 3$ yields a group of transformations $\langle \sigma_1, \sigma_2, \sigma_3, \tau_1, \tau_2, \tau_3 \rangle$ which, it can be verified, is the transformation group for initial data configurations of the 5-simplex pair-configuration of Proposition 5. In this case the variables x_1, x_2, x_3 and y_1, y_2, y_3 of (7.1) are associated with a pair of disjoint duad-syntheses (cf. Definition 5) of the 5-simplex (despite




Fig. 10. Coxeter-Dynkin diagram for the index-2 normal subgroup of the vertex-map group.

this we retain the phrase *vertex maps*, which recalls the original context of the construction). Thus combining the vertex maps provides a constructive method to recover the 5-simplex domain that was obtained in Section 6. On this basis, we now investigate the group generated by the combined vertex maps for generic n .

Analysis of this group is simplified by the observation that vertex maps from the two classes can be related,

$$\tau_j = \sigma_j \omega \sigma_j, \quad (7.14)$$

by introducing the map ω that interchanges columns

$$\omega := \begin{bmatrix} x_1, y_1 \\ \vdots \\ x_n, y_n \end{bmatrix} \mapsto \begin{bmatrix} y_1, x_1 \\ \vdots \\ y_n, x_n \end{bmatrix}. \quad (7.15)$$

We assume the integer n is fixed and consider the group generated by

$$\sigma_1, \dots, \sigma_m, \omega, \quad (7.16)$$

for $m < n$. This has the advantage that if $m' < m$, then $\langle \sigma_1, \dots, \sigma_{m'}, \omega \rangle$ is a subgroup of $\langle \sigma_1, \dots, \sigma_m, \omega \rangle$. The following relations are key:

$$\begin{aligned} \sigma_i^2 &= \omega^2 = (\sigma_i \omega)^4 = \text{id}, \\ (\sigma_i \sigma_j)^2 &= (\sigma_i \omega \sigma_j \omega)^3 = \text{id}, & |\{i, j\}| &= 2, \\ ((\sigma_i \sigma_j \omega)^2 \sigma_k \omega)^2 &= \text{id}, & |\{i, j, k\}| &= 3, \end{aligned} \quad (7.17)$$

they encode the stronger 5-simplex consistency property.

The main result of this section is the characterisation of this group for generic $m < n$.

Proposition 6. *The finitely presented group with generators (7.16) and relations (7.17) contains an index-2 normal subgroup isomorphic to the Coxeter group of Figure 10.*

Proof. The index-2 normal subgroup can be written as

$$\langle \omega, \sigma_1 \omega \sigma_1, \sigma_1 \sigma_2, \sigma_1 \sigma_3, \dots, \sigma_1 \sigma_m \rangle. \quad (7.18)$$

Group	Order	Isomorphic to
$\langle \sigma_1, \sigma_2, \omega \rangle$	2×36	$A_1 \ltimes (A_2 \times A_2)$
$\langle \sigma_1, \sigma_2, \sigma_3, \omega \rangle$	2×720	$A_1 \ltimes A_5$
$\langle \sigma_1, \sigma_2, \sigma_3, \sigma_4, \omega \rangle$	2×51840	$A_1 \ltimes E_6$
$\langle \sigma_1, \sigma_2, \sigma_3, \sigma_4, \sigma_5, \omega \rangle$	$2 \times \infty$	$A_1 \ltimes \tilde{E}_6$

Table 2. Sequence of subgroups of the complete vertex-map group.

The claim is that this group is isomorphic to the group represented in Figure 10. We proceed by exhibiting an isomorphism:

$$\begin{aligned}
t_1 &= \omega, & \omega &= t_1, \\
t_2 &= \sigma_1 \omega \sigma_1, & \sigma_1 \omega \sigma_1 &= t_2, \\
t_3 &= \omega \sigma_1 \sigma_2 \omega \sigma_2 \sigma_1 \omega, & \sigma_1 \sigma_2 &= t_3 t_4, \\
t_4 &= (\omega \sigma_1 \sigma_2)^3, & \sigma_1 \sigma_3 &= (t_3 t_4)^{t_5}, \\
t_5 &= (\omega \sigma_2 \sigma_3)^3, & \sigma_1 \sigma_4 &= (t_3 t_4)^{t_5 t_6}, \\
&\vdots & &\vdots \\
t_{m+2} &= (\omega \sigma_{m-1} \sigma_m)^3, & \sigma_1 \sigma_m &= (t_3 t_4)^{t_5 t_6 \cdots t_{m+2}}.
\end{aligned} \tag{7.19}$$

Here the notation f^g denotes conjugation of f by g , $f^g = g^{-1}fg$. The group elements $t_1, \dots, t_{m+2}$ are associated with the labelled nodes of Figure 10. An advantage of the chosen isomorphism is that to verify it, calculation is required only up to the case $m = 4$, because beyond this the new relations which appear are similar to existing ones that have already been checked. This can be seen by inspection of the isomorphism (7.19), the relations (7.17) and the Figure 10. At the value $m = 4$ the groups in question are finite and the isomorphism may be checked using computation based on a permutation representation. The computer algebra system Magma [33] was used to construct an isomorphism at $m = 4$.

Note that mappings of the form $(\omega \sigma_i \sigma_{i+1})^3$, which appear in (7.19), correspond to pure permutations in the vertex-map group, specifically transposition of rows i and $i + 1$ of the array (7.1). In particular, we note that for $i \in \{1, \dots, m - 1\}$, $t_{i+3} = (\omega \sigma_i \sigma_{i+1})^3 = s_i^\rho$, where $s_1, \dots, s_{m-1}$ are the braid-type mappings.

We also remark on one subtlety, namely that $m < n$ is required for the relations (7.17) to be complete. In fact, due to the groups being finite, it is easy to check that the following holds.

Lemma 2. *If $m = n \in \{3, 4\}$ then $\langle \sigma_1, \dots, \sigma_n, \omega \rangle = \langle \omega, \sigma_1 \omega \sigma_1, \sigma_1 \sigma_2, \dots, \sigma_1 \sigma_n \rangle$.*

This would contradict Proposition 6 if the relations were complete in the case $m = n$. It may continue to hold for all $m = n > 2$ (but is not true if $m = n = 2$).

The group relations (7.17) and Proposition 6 can be seen as a fairly complete characterisation of the algebraic structure associated with the family of idempotent biquadratics that share a common orbit. To understand what this means for the associated discrete dynamics, the group structure requires some further interpretation. Within Figure 10 one can see the symmetry group of a single triplet-pair corresponding to $m = 2$, and of the 5-simplex pair-configuration when $m = 3$. Continuing, the combined vertex-map group decomposes as in Table 2. In particular, that the group becomes infinite beyond $m = 4$ represents

a significant and un-anticipated depth beyond the kind of lattice geometry of the oriented n -simplex or the n -dimensional hypercube. A natural step towards understanding the dynamics of the ambient space as a whole, is to relate the lattice geometry to the group.

8. The generalised lattice geometry

The abstract structure of the vertex-map group encodes the full ambient domain on which it is natural to consider model (4.4). Here we explain how the geometry of the domain may be decoded from the group, make contact with the configuration of the 27 lines of a cubic surface in the case $n = 4$, and discuss the dynamics associated with subgroups.

8.1. Decoding the geometry from the group. Recall from Section 6 that in general the domain of (4.4) is a set of variables arranged into triplet-pairs (cf. Definition 2). Consider now the following associations,

$$\begin{aligned} \text{variables} &\leftrightarrow V := \{\sigma_1^g : g \in \langle \sigma_1, \dots, \sigma_n, \omega \rangle\}, \\ \text{triplet-pairs} &\leftrightarrow T := \{\langle \sigma_1, \sigma_2, \omega \rangle^g : g \in \langle \sigma_1, \dots, \sigma_n, \omega \rangle\}. \end{aligned} \quad (8.1)$$

Again here f^g denotes conjugation of f by g , $f^g = g^{-1}fg$. Thus a set of variables V and triplet-pairs T are identified with particular conjugacy classes of the vertex map group. It may be decided whether a variable is contained within a triplet-pair in the natural way: the triplet-pair corresponding to $\pi \in T$ contains the variable corresponding to $\sigma \in V$ when $\sigma \in \pi$. In particular, the internal structure of the triplet-pair corresponding to $\pi \in T$ may be recovered by taking the intersection $V \cap \pi$ and considering how the action of π permutes these elements of V .

Note that all of $\sigma_1, \dots, \sigma_n$ are conjugate (this can be seen from the definition of σ_j (7.5), (7.6) by considering the pure-permutation sub-group), therefore all are identified with variables. The vertex-type initial data can also be recovered from the group, it corresponds to variables associated with the set

$$\{\sigma_1, \dots, \sigma_n, \sigma_1^\omega, \dots, \sigma_n^\omega\}, \quad (8.2)$$

or any set conjugate to this one. Furthermore, arranging this set in the form

$$\begin{bmatrix} \sigma_1^\omega, \sigma_1 \\ \vdots \\ \sigma_n^\omega, \sigma_n \end{bmatrix} \quad (8.3)$$

provides a method to see the consistency of identifications (8.1). One checks that the natural action of the group on (8.3) (i.e., by conjugation of each component), is consistent with its default action on (7.1). It suffices to check the action of the generators.

The combinatorial considerations raised in Section 6 are also taken care of automatically by this construction.

Proposition 7. *The variables and triplet-pairs defined by associations (8.1) form a symmetry-complete pair-configuration (cf. Definitions 3 and 4).*

Proof. The domain is clearly a pair-configuration because all members of V and T are on an equal footing amongst themselves (are conjugate). Now observe that the symmetry group of the triplet-pair associated with $\pi \in T$ can be identified with the inner automorphism group of π . Clearly the action of π extends naturally to the whole group, and thus the pair-configuration is symmetry-complete.

8.2. Case $n = 4$. The case $n = 4$ is significant, being the last finite case in the sequence of complete vertex-map groups, cf. Table 2. This group, which is isomorphic to E_6 when $n = m = 4$ (Table 2 shows $n > m = 4$, cf. Lemma 2), was studied by Jorden [34] as the group of incidence-preserving permutations of the 27 straight lines on a cubic surface, which he connected with the trisection of periods of quadruply-periodic functions. Subsequently Burkhardt [35], through the theta-function approach introduced by Jordan, obtained a linear representation of the group, and in particular an invariant of the representation that, when set equal to zero, defined a quartic hypersurface (quartic primal). Amongst the substantial literature on this subject, we cite here mainly the work of Baker [36] who studied the geometry of Burkhardt's quartic. This study was purely in terms of position in four-dimensional space of the 45 singular points, called nodes, which he found were in natural correspondence with 45 particular elements of the group, called projections. The set of projections corresponds in turn to the set V in (8.1), i.e., the variables of the $n = 4$ pair-configuration. The projections were proven to form a complete conjugacy class by Todd [37], who also makes a concise summary of Bakers extensive geometric description of the 45 nodes.

We recount a part of this geometry in order to identify in Bakers scheme what we have called triplet-pairs here. The 45 nodes lie by threes on lines, called κ -lines, with 16 lines through each node and 240 lines in total. The nodes and κ -lines therefore form a $(45_{16}240_3)$ configuration. The set of all κ -lines intersecting a particular one contain every node of the configuration except three. Those three remaining nodes themselves lie on a second κ -line, which is called polar to the first. The relationship is reciprocal, and the 120 pairs of polar κ -lines can be identified with the triplet-pairs T in (8.1).

Within the $n = 4$ pair-configuration there are 36 distinct copies of the $n = 3$ (5-simplex) pair-configuration. This sub-configuration is left unnamed by Baker, but is called a 15-set by Todd. On the other hand, one of Bakers additional contributions was to provide a coordinatization, and this implicitly acknowledges the 15-set because it directly extends the natural coordinatization of it given earlier in Proposition 4. It allows to write down the $n = 4$ pair-configuration as follows.

Proposition 8 (notation due to Baker [36]). *The pair-configuration defined by associations (8.1) in the case $n = 4$ consists of 45 variables arranged into 120 triplet-pairs. If we denote 15 of the variables by*

$$x_{ij} = x_{ji}, \quad i, j \in \{1, 2, 3, 4, 5, 6\}, |\{i, j\}| = 2, \quad (8.4)$$

and 30 of them by

$$x_{(ij,kl,mn)} = x_{(ji,kl,mn)} = x_{(kl,mn,ij)}, \quad \{i, j, k, l, m, n\} = \{1, 2, 3, 4, 5, 6\}, \quad (8.5)$$

then the 120 triplet-pairs may be written as follows: there are 10 of the form

$$x_{ij}, x_{jk}, x_{ki}, \quad x_{lm}, x_{mn}, x_{nl}, \quad (8.6)$$

90 of the form

$$x_{ij}, x_{(ik,jl,mn)}, x_{(jk,il,mn)}, \quad x_{mn}, x_{(mk,nl,ij)}, x_{(nk,ml,ij)}, \quad (8.7)$$

and 20 of the form

$$x_{(ij,kl,mn)}, x_{(il,kn,mj)}, x_{(in,kj,ml)}, \quad x_{(ij,mn,kl)}, x_{(il,mj,kn)}, x_{(in,ml,kj)}, \quad (8.8)$$

where in (8.6), (8.7) and (8.8), $\{i, j, k, l, m, n\} = \{1, 2, 3, 4, 5, 6\}$. The following set of variables may be given as initial data for this pair-configuration

$$\begin{bmatrix} x_{12} & , & x_{54} \\ x_{34} & , & x_{16} \\ x_{56} & , & x_{32} \\ x_{(12,34,56)} & , & x_{(16,54,32)} \end{bmatrix}. \quad (8.9)$$

The configuration also carries in total $540 = 36 \times 15$ copies of model (6.5): the pairwise intersection of the 36 copies of the 5-simplex sub-configuration is of two kinds, either a triplet-pair, or a duad-syntheme, thus no two share the same edge-tetrahedron pair of Figure 9. We also remark there are in total 135 distinct copies of the 4-cube, which is the quad-graph domain generalised by this configuration.

8.3. Subgroup dynamics and integrability. By construction, the complete vertex-map group contains A_n and BC_n subgroups corresponding to the n -simplex and n -cube domains. In both of these domains it is envisaged that n is arbitrarily large, and that the number of degrees of freedom is therefore un-bounded. Integrability on the quad-graph domain is of KdV-type, whereas the integrability on the triangle-graph domain (discussed in Section 5) is of a simpler kind.

Due to the nature of the complete vertex map group, the unbounded domain required for non-trivial dynamics is also present for finite n , the simplest case being $n = 5$. The group in this case is the affine reflection group $\tilde{E}_6$, such affine groups are familiar from dynamics in the Painlevé setting, though the birational representation here looks very different. Nevertheless it would be natural to adopt the similar point of view, that dynamics are associated with the normal abelian subgroup, whilst the action of the finite quotient group (E_6) corresponds to symmetries of the dynamical system. This would give a natural notion of integrability, but it remains open whether the systems here can actually be integrated in terms of (discrete) Painlevé transcendents or their generalisations.

Restriction to subgroup dynamics by taking a reduced set of generators leads to an additional variety of possibilities. An interesting case would be the group

$$\langle t_1, t_3, \dots, t_{10} \rangle, \quad (8.10)$$

written in terms of elements given in (7.19), which constitutes a birational representation of $\tilde{E}_8$. Note that when restricting to a proper subgroup the associations (8.1) will in general require modification.

Due to the rich structure of the general vertex map group, it is natural to expect that subgroup dynamics more generally can also be considered integrable in some sense. This is something which could potentially be tested by the notion of algebraic entropy [38, 39] for instance.

9. Discussion

Non-metric domains. It is relevant here to address the fact that the ambient domain, a symmetry-complete pair-configuration (Definitions 3 and 4) a-priori carries no natural notion of distance, we loosely refer to such a domain as being non-metric. To use the language of dynamical systems can begin to be awkward in such situations. On the other hand, this domain is encountered here as the natural destination along the path of the consistency framework for integrable systems. What is most striking is the inherent potential of such domains to unify integrable systems from quite different settings. In particular, the one model here encompasses dynamics associated with a quad-graph, a triangle-graph, affine reflection groups and more (cf. Section 8.3).

Symmetry-completeness and consistency. Here we have identified the consistency configuration underlying the integrability of two quad-graph models as being in fact a sub-configuration of a larger whole. Aside from the algebraic motivation for identifying this generalised picture, which is based on the idempotent biquadratic and therefore specific to the particular models at hand, it is shown that it can also be discovered through the perfectly transferable and wholly combinatorial concept of symmetrization (cf. Definition 4 and Proposition 4). Moreover, the technique based on vertex maps (more generally, mappings between initial data sets of the configuration) provides a method to obtain the symmetrized configuration. These methods consider the consistency of itself and therefore have application beyond the system studied here.

Geometry. Adler, Bobenko and Suris, when introducing the scalar quadrirational models [2], gave also a geometric incidence picture in terms of pencils of conics. The refinement of this geometric picture is potentially very interesting. In particular offering the possibility to make a connection with the configuration of 27 lines on the cubic surface beyond the purely combinatorial relationship given.

Generalisations. Underlying the system studied here are the idempotent biquadratics. More specifically, the family that share a common fixed orbit, that we have denoted $z_0, z_1, z_2 \in \mathbb{C} \cup \{\infty\}$ throughout. The invariance under permutation of orbits raises the question of whether there is a yet more general consistency configuration in which this fixed orbit participates on the same level as all the others.

Beyond the idempotent biquadratic is the idempotent bicubic. This is defined by the similar Bezout-type expression (3.6), but with r_1 and r_2 now quartic polynomials. As a correspondence, it directly generalises the biquadratic case, and the three-orbit constraint is again rational. This is therefore a potentially significant object and will be the subject of future work.

A. Idempotent biquadratic polynomials in the Weierstrass theory of elliptic functions

Consider the symmetric biquadratic polynomial,

$$h(x, y) = c_0 + c_1(x + y) + c_2xy + c_3(x^2 + y^2) + c_4xy(x + y) + c_5x^2y^2, \quad (\text{A.1})$$

$c_0, \dots, c_5 \in \mathbb{C}$. As mentioned in the main text, its orbits (3.2), (3.3) may be taken as a definition of elliptic functions in the discrete setting. We develop here an aspect of this discrete theory connected to the Weierstrass addition formula, and relate it to the idempotent class of biquadratics.

A basic role in this aspect of the Weierstrass theory is played by the discriminant polynomial r associated with h , which is defined as

$$\Delta[h(x, y), y] = r(x), \quad \Delta := \text{discriminant}. \quad (\text{A.2})$$

This terminology is unambiguous due to the symmetry of h . Henceforth we assume that r has at least one simple root. The fundamental property on the level of biquadratics is a kind of closure under composition within the family with shared discriminant polynomial. More precisely, if biquadratics h_1 and h_2 have, up to a scalar multiple, a common discriminant polynomial r , then the composition of h_1 and h_2 takes the form

$$\text{res}[h_1(x, y), h_2(y, z), y] = h_3(x, z)h_4(x, z), \quad \text{res} := \text{resultant}, \quad (\text{A.3})$$

where h_3 and h_4 are again symmetric biquadratic polynomials whose discriminant polynomial is a (now possibly zero) scalar multiple of r . The appearance of two factors on the right-hand-side of (A.3) means this composition gives a structure which is sometimes called a two-valued group.

A natural parameterisation of this group is obtained through the evaluation map,

$$h \mapsto \alpha : h(e_0, \alpha) = 0. \quad (\text{A.4})$$

Choosing the point of evaluation, $e_0 \in \mathbb{C} \cup \{\infty\}$, to be the assumed simple root of r , makes the evaluation map single-valued. Furthermore, h can be reconstructed (up to constant scalar multiple) from its discriminant polynomial and the associated parameter α . Specifically

$$h(x, y) \sim H(x, y, \alpha) \quad (\text{A.5})$$

where H is a triquadratic polynomial which may be expressed as a discriminant

$$H := \Delta \left[\frac{r_1(t)r_2(e_0) - r_2(t)r_1(e_0)}{t - e_0}, t \right], \quad (\text{A.6})$$

where t is a dummy variable, and

$$r_1(t) \sim \frac{r(t)}{t - e_0}, \quad r_2(t) = (t - x)(t - y)(t - \alpha). \quad (\text{A.7})$$

Observe that H is invariant under permutations of the three variables x, y and α . This permutation symmetry has the remarkable consequence that the parameters $\alpha_1, \dots, \alpha_4$ associated with the biquadratics appearing in the composition formula (A.3),

$$h_i \mapsto \alpha_i : h_i(e_0, \alpha_i) = 0 \quad \Rightarrow \quad h_i(x, y) \sim H(x, y, \alpha_i), \quad i \in \{1, 2, 3, 4\}, \quad (\text{A.8})$$

are themselves related by the same polynomial H ,

$$H(\alpha_1, \alpha_2, z) = 0 \Leftrightarrow z \in \{\alpha_3, \alpha_4\}. \quad (\text{A.9})$$

In particular, the binary operation $(x, y) \mapsto z$ defined by equation $H(x, y, z) = 0$ endows $\mathbb{C} \cup \{\infty\}$ with the structure of an abelian two-valued group. Most of the important features of H (A.6) can be seen from its canonical form, the case $e_0 = \infty$, $r(t) = 4t^3 - g_2t - g_3$:

$$H(x, y, z) = (xy + yz + zx + \frac{1}{4}g_2)^2 - 4(x + y + z)(xyz - \frac{1}{4}g_3). \quad (\text{A.10})$$

Those features being that it appears as the discriminant of a polynomial which is degree-one in each of three variables, that it is symmetric in the three variables, and, through the Weierstrass addition formula

$$H(\wp(\alpha), \wp(\beta), \wp(\gamma)) = 0 \Leftrightarrow \wp(\gamma) = \wp(\alpha \pm \beta), \quad (\text{A.11})$$

also that it defines a two-valued abelian group structure. The more generic form (A.6) reveals the connection with the Bezout-type expression for the idempotent biquadratic correspondences (3.6).

Acknowledgements. This research was funded by Australian Research Council Discovery Grant DP 110104151. Thanks are due to Pavlos Kassotakis for many discussions, and David Howden for advice on Magma.

References

1. J. Atkinson and M. Nieszporski. Multi-quadratic quad equations: integrable cases from a factorised discriminant hypothesis. *Intl. Math. Res. Not.*, 2012. submitted.
2. V. E. Adler, A. I. Bobenko, and Yu. B. Suris. Geometry of Yang-Baxter maps: pencils of conics and quadrirational mappings. *Comm. Anal. Geom.*, 12(5):967–1007, 2004.
3. H. D. Wahlquist and F. B. Estabrook. Bäcklund transformation for solutions of the Korteweg-de Vries equation. *Phys. Rev. Lett.*, 31:1386–1390, 1973.
4. R. Hirota. Nonlinear partial difference equation I: A difference analogue of the KdV equation. *J. Phys. Soc. Jp.*, 43:1429–1433, 1977.
5. F. W. Nijhoff, G. R. W. Quispel, and H. W. Capel. Direct linearization of nonlinear difference equations. *Phys. Lett. A*, 97:125–128, 1983.
6. F. W. Nijhoff and H. W. Capel. The discrete Korteweg-de Vries equation. *Acta Appl. Math.*, 39(1-3):133–158, 1995.
7. V. E. Adler, A. I. Bobenko, and Yu. B. Suris. Classification of integrable equations on quad-graphs. The consistency approach. *Comm. Math. Phys.*, 233(3):513–543, 2003.
8. V. E. Adler, A. I. Bobenko, and Yu. B. Suris. Discrete nonlinear hyperbolic equations. Classification of integrable cases. *Funct. Anal. Appl.*, 43(1):3–17, 2009.
9. J. Atkinson. Bäcklund transformations for integrable lattice equations. *J. Phys. A: Math. Theor.*, 41:Art. No. 135202, 2008.
10. V. G. Papageorgiou, A. G. Tongas, and A. P. Veselov. Yang-Baxter maps and symmetries of integrable equations on quad-graphs. *J. Math. Phys.*, 47:Art. no. 083502, 2006.
11. V. G. Papageorgiou, Yu. B. Suris, A. G. Tongas, and A. P. Veselov. On quadrirational Yang-Baxter maps. *SIGMA* 6, 033:9pp, 2010.
12. P. Kassotakis and M. Nieszporski. On non-multiaffine consistent around the cube lattice equations. *Physics Letters A*, 376:3135–3140, 2012.
13. P. Kassotakis and M. Nieszporski. Families of integrable equations. *SIGMA*, 7:100, 2011.
14. J. Atkinson and N. Joshi. The Schwarzian variable associated with discrete KdV-type equations. *Nonlinearity*, 25(6):1851–1866, 2010.
15. J. Atkinson. A multidimensionally consistent version of Hirota’s discrete KdV equation. *J. Phys. A: Math. Theor.*, 45:Art. No. 222001, 2012.

16. A. P. Veselov. Yang-Baxter maps: dynamical point of view. *Math. Soc. Japan memoirs*, 17, Combinatorial aspects of Integrable Systems:pp167, 2006.
17. M. Noumi and Y. Yamada. Affine weyl groups, discrete dynamical systems and Painlevé equations. *Communications in Mathematical Physics*, 199(2):281–295, 1998.
18. H. Sakai. Rational surfaces associated with affine root systems and geometry of the painlevé equations. *Communications in Mathematical Physics*, 220(1):165–229, 2001.
19. K. Kajiwara, M. Noumi, and Y. Yamada. Discrete dynamical systems with $W(A_{m-1}^{(1)} \times A_{n-1}^{(1)})$ symmetry. *Letters in Mathematical Physics*, 60(3):211–219, 2002.
20. K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta, and Y. Yamada. ${}_{10}E_9$ solution to the elliptic Painlevé equation. *J. Phys. A: Math. Gen.*, 36:L263–L272, 2003.
21. T. Tsuda. A geometric approach to tau-functions of difference Painlevé equations. *Letters in Mathematical Physics*, 85(1):65–78, 2008.
22. K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta, and Y. Yamada. Point configurations, Cremona transformations and the elliptic difference Painlevé equation. *Séminaires et Congrès*, 14.
23. A. I. Bobenko and Yu. B. Suris. Integrable systems on quad-graphs. *Intl. Math. Res. Notices*, 11:573–611, 2002.
24. V. E. Adler and A. P. Veselov. Cauchy problem for integrable discrete equations on quad-graphs. *Acta Appl. Math.*, 84(2):237–262, 2004.
25. P. H. van-der Kamp. Initial value problems for quad equations. *arXiv:1210.1414 [nlin.SI]*, 2012.
26. J. Atkinson. Integrable lattice equations: Connection to the Möbius group, Bäcklund transformations and solutions. *PhD thesis, The University of Leeds*, 2008.
27. I. M. Gelfand, M. M. Kapranov, and A. V. Zelevinsky. *Discriminants, resultants and multidimensional determinants*. Mathematics: Theory and Applications. Birkhäuser, 1994.
28. H. Hancock. *Lectures on the theory of elliptic functions*. New York: John Wiley & sons, London: Chapman and Hall, 1910.
29. J. J. Sylvester. Elementary researches in the analysis of combinatorial aggregation. *Philosophical Magazine*, XXIV:285–296, 1844.
30. H. S. M. Coxeter. Self-dual configurations and regular graphs. *Bulletin of the American Mathematical Society*, 56:413–455, 1950.
31. A. P. Veselov. Yang-Baxter maps and integrable dynamics. *Phys. Lett. A*, 314:214–221, 2003.
32. T. E. Kouloukas and V. G. Papageorgiou. Entwining Yang-Baxter maps and integrable lattices. *Banach center publ.*, 93:163–175, 2010.
33. W. Bosma, J. Cannon, and C. Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).
34. C. Jordan. *Traité des substitutions et des équations algébriques*. Paris: Gauthier-Villars, 1870.
35. H. Burkhardt. Untersuchungen aus dem gebiete der hyperelliptischen modulfunktionen erster theil. *Mathematische Annalen*, 36(3):371–434, 1890.
36. H. F. Baker. *A locus with 25920 linear self-transformations*, volume 39 of *Cambridge tracts in mathematics and mathematical physics*. Cambridge University Press, 1946.
37. J. A. Todd. On the simple group of order 25920. *Proc. R. Soc. Lond. A*, 189:326–358, 1947.
38. G. Falqui and C. M. Viallet. Singularity, complexity, and quasi-integrability of rational mappings. *Commun. Math. Phys.*, 154:111–125, 1993.
39. M. P. Bellon and C.-M. Viallet. Algebraic entropy. *Communications in Mathematical Physics*, 204(2):425–437, 1999.